



MedStar Health
Institute for Quality and Safety

MedStarBloodless.org

International Training Center for
Bloodless Medicine and Surgery

MedStar Franklin Square Medical Center
MedStar Georgetown University Hospital
MedStar Good Samaritan Hospital
MedStar Harbor Hospital
MedStar Montgomery Medical Center
MedStar National Rehabilitation Network
MedStar Southern Maryland Hospital Center
MedStar St. Mary's Hospital
MedStar Union Memorial Hospital
MedStar Washington Hospital Center
MedStar Family Choice
MedStar Medical Group
MedStar PromptCare
MedStar Visiting Nurse Association
MedStar Institute for Innovation
MedStar Health Research Institute

CONTRIBUTED REPORTS

**Total Hip Replacement
without Blood Transfusion:
53-Year-Old Male with
Cardiac Arrhythmia**

Timothy Penn, MD - December 3, 2015

Knowledge and Compassion
Focused on You

Total Hip Replacement without Blood Transfusion: 53-Year-Old Male with Cardiac Arrhythmia

Timothy Penn, MD

Abstract

Perioperative blood loss is a significant concern for patients undergoing total joint arthroplasty. A growing body of evidence has shown tranexamic acid (TXA) to be effective in decreasing perioperative blood loss and transfusion requirements in both primary and revision hip and knee arthroplasty. Dr. Penn reports using TXA in successful hip replacement in a patient with cardiac arrhythmia.

Case Report

A 53-year-old male Jehovah's Witness presented with severe osteoarthritis of the left hip. He had long-term non-operative management including exercise, NSAIDs, and intra-articular injections. Ultimately, these didn't provide lasting relief and he opted for hip replacement. Medical issues included cardiac arrhythmia, chronic back issues, hypertension and elevated cholesterol. His pre-operative Hgb was 16.7. In December 2015 he underwent a left total hip replacement through an anterior approach. Perioperative management included normothermia, closed loop cell saver, and 1 gram tranexamic acid administered IV one hour pre-incision, with intraoperative administration as a direct application into the surgical wound right before closure. Intra-op blood loss was measured to be 500 ml, with 400 ml returned through the closed loop cell saver. Post-operatively his Hgb was 15.4 in the recovery room, dropping to 14.1, noted a week later when he was seen in the ER for opiate induced constipation. 2 weeks post-operatively, he was walking with a cane. At 2 months his issues were all back related, and the left hip was essentially pain free. He went on to have back surgery 5 months after his hip surgery. He was seen for routine

x-ray follow-up at 1 and 2 years post-surgery and continues to do well. Next x-rays are scheduled for the 5-year point from his surgery.

Dr. Penn's Notes

The TXA comes as 1 gram in a 10 mL vial. The pre-operative dose is given within the hour before incision, 1 gram IV over 10 minutes or more. Sometimes it is given full strength, but some of the nurses will dilute it with 10 mL of normal saline, doubling the volume. The reason it is given slowly is because it can cause nausea if given too fast. The exact timing of the pre-operative administration isn't critical.

The topical is undiluted 1 gram placed into the wound at closure. An additional gram around 3 hours post-OR was tried but seemed to cause a leukocytosis with some consistency. The third dose didn't seem to make a difference with respect to the postoperative Hgb. No special consideration is given for presence of stents or anything else, as this material hasn't been found to be thrombogenic. There is evidence that oral preoperative dosing is as effective as IV, but since our patients are NPO, we opt for the IV.

About the Author



Timothy Penn, M.D.
Orthopedic Surgery,
Orthopedics

Disclaimer

This content is provided for general informational purposes only and is not intended as, nor should it be considered a substitute for, professional medical advice. Do not use the information on this website for diagnosing or treating any medical or health condition. If you have or suspect you have a medical problem, promptly contact your professional healthcare provider.